

Erratum: Light-induced desorption of alkali-metal atoms from paraffin coating [Phys. Rev. A **66**, 042903 (2002)]

E. B. Alexandrov and M. V. Balabas
S. I. Vavilov State Optical Institute, St. Petersburg, 199034 Russia

D. Budker*
*Department of Physics, University of California at Berkeley, Berkeley, California 94720-7300 and
Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720*

D. English, D. F. Kimball, C.-H. Li, and V. V. Yashchuk
Department of Physics, University of California at Berkeley, Berkeley, California 94720-7300
(Dated: February 2, 2008)

PACS numbers: PACS. 34.50.Dy, 79.20.La

In the above manuscript, there was an error in the numeric computation [based on Eqs. (6) and (7)] predicting the time-dependent alkali vapor density $n(t)$ and number of alkali atoms in the paraffin coating $N_c(t)$. Once the error in the computation was remedied, it was necessary to adjust two parameters [ρ (probability for adsorption of atoms by the paraffin coating per wall collision) and ξ (exchange rate between atoms in the stem of the cell and the volume of the cell)] in order to obtain the same agreement between the theory and the data as seen in Figs. 9 and 10. This modifies the results for ρ and ξ listed in Table II as follows:

TABLE II: Average values of various light-independent parameters from the model extracted from fits to the data from Secs. III B, III C, and III D.

Parameter	Type (a) cell	Type (b) cell
Γ	$5(2) \times 10^{-4} \text{ s}^{-1}$	$7(4) \times 10^{-4} \text{ s}^{-1}$
γ_d	$2(1) \times 10^{-4} \text{ s}^{-1}$	$2(1) \times 10^{-4} \text{ s}^{-1}$
ρ	$4.9(3) \times 10^{-6}$	$4(2) \times 10^{-6}$
ξ	$5.0(2) \text{ cm}^3/\text{s}$	$1.49(3) \text{ cm}^3/\text{s}$

Therefore the average over all data for ρ should be $4(1) \times 10^{-6}$. This in fact improves agreement with the values of ρ determined in other work [1, 2, 3, 4], in which ρ was found to be between $\sim 10^{-6}$ and $\sim 3 \times 10^{-6}$. This in turn means that the number of atoms estimated to be in the coating N_c should be larger by a factor of about 30 [see, for example, Eq. (9)]. This would correspondingly change the scale of the lower plots in Figs. 9 and 10.

The adjustment to the value for ξ changes our estimate of the number of atoms adsorbed by the coating during the “ripening” process to $\sim 10^{17}$, but this in no way changes our conclusion that the number of atoms available for desorption is considerably smaller than the number of atoms absorbed during cell ripening.

We are deeply indebted to A. I. Okunevich for bringing this mistake to our attention.

-
- [1] M. A. Bouchiat and J. Brossel, Phys. Rev. **147**, 41 (1966); M. A. Bouchiat, Ph. D. Thesis, University of Paris, 1964.
 - [2] V. Liberman and R. J. Knize, Phys. Rev. A **34**, 5115 (1986).
 - [3] M. V. Balabas and V. A. Bonch-Bruevich, Pis'ma Zh. Tekh. Fiz. **19**, 6 (1993) [Tech. Phys. Lett. **19**, 191 (1993)].
 - [4] M. V. Balabas and S. G. Przhibel'skii, Khimicheskaya Fizika **14**, 125 (1995).

*Electronic address: budker@socrates.berkeley.edu